

RAISE THE EMOTIONAL THRESHOLD

Smart young Americans are changing *today*

_WHY DON'T YOU? You'll be glad tomorrow . . .



Warning: excessive ingestion of this substance has been found to cause: confusion, brain damage, nausea, drowsiness, sudden change in mood or personality, seizures, headache, memory loss, restlessness, irritability, muscle pain or weakness, death, coma, respiratory arrest, lethargy, fatigue, cramping, dizziness



~~IIII~~ ~~IIII~~
~~IIII~~ ~~IIII~~ III

xanax morphine adderall klonopin valium ambien heroin fentanyl codeine vicodin dilaudid oxy-contin ritalin methadone ativan opana vyvanse phenobarbital dextromethorphan concerta tu-inal lortab demerol suboxone roxicet dol-abstral s-lunesta biphetamine seconal duragesic prozac es-paroxetine rtraline fluvoxamine

percocet numorphan trazodone ophine sublimaze eroquel librium nembutal dexedrine oxycodone citalopram celexa se-vilazodone

cocaine loperamide hydromorphone avinza tramadol amitriptylin ketamine lexapro cymbalta wellb-utrin meperidine temazepam suvorexant triaz-olam estazolam flurazepam clorazepate prosom carfentanil exalgo quazepam versed fluoxetine

THE GREAT PAIN CURE.

Cautions and Side Effects: At daytime sedative dosage levels, drowsiness, skin rash, "hangover" and systemic disturbances are seldom seen. To be used with caution in moderate to severe hepatic disease. If a hypersensitivity reaction should occur, discontinue use. (*Warning:* May be habit forming.)

Dosage: As a daytime sedative, 15 mg. (¼ gr.) to 30 mg. (½ gr.) t.i.d. or q.i.d.

Available: Tablets, 15 mg. (¼ gr.), 30 mg. (½ gr.), 50 mg. (¾ gr.), 100 mg. (1½ gr.); Elixir [30 mg. (½ gr.)/5 cc.] Alcohol 7%.

RUN! LITTLE COCKROACH RUN!

a new standard in tranquilization